

フェビモ® CHIBISHIMO (Ointment for frostbite)

Ointment for frostbite.

Class 2 drug



Effective for frostbite, itchiness, improves circulation to help bring skin close to a normal state.

An easy-to-use heat type ointment for children to adults.

MADE IN JAPAN

Indications

frostbite, itchiness, eczema, and dermatitis

Dosage and Administration

Apply an appropriate amount several times a day to the affected area.

(Precautions related to Dosage and Administration)

- (1) When used for children, please use under the guidance and supervision of the guardian.
- (2) Do not touch around the eyes or around the mouth with the ointment on your hands.
- (3) Keep out of eyes. If by chance the product gets in the eyes, rinse immediately with water or tepid water. Furthermore, when symptoms are severe, please receive medical attention from an ophthalmologist.
- (4) For external use only.

Ingredient, content (per 100g), and action

Ingredient	Content	Action
diphenhydramine hydrochloride	2.0g	Blocks histaminic activity, and inhibits itchiness well.
lidocain	2.0g	Provides fast relief of itchiness by way of local anesthesia action.
glycyrrhetinate	0.5g	Inhibits inflammation such as chillblains and eczema.
tocopherol acetate	0.1g	Dilates peripheral blood vessels to improve circulation of the affected area.

Contains macrogol and polysorbate 80 as the additive.

※This product does not contain steroid components.

Precautions

Things to avoid

Things to avoid

(if not respected, the present symptoms will exacerbate, and side effects will readily develop)

1. Do not use the ointment in the following areas:

Around the eyes, and mucous membranes, etc.

Things to consult

1. Before using the product, the following individuals should consult their doctor, pharmacist, or registered sales clerk:

- (1) Individuals receiving medical treatment from a doctor.

(2) Individuals who have a history of drug-induced allergic reaction.

(3) Individuals with severe dampness or inflammation.

2. After usage, if any of the following symptoms occur, they could be side effects; thus, discontinue usage of the ointment immediately, and please consult your doctor, pharmacist, or registered sales clerk with this document.

Site concerned	Symptoms
Skin	Eruption, redness, itchiness, and swelling

3. If symptoms do not improve despite usage for 5 - 6 days, discontinue usage, and please consult your doctor, pharmacist, or registered sales clerk with this document.