

コンジスイQ

KONJISUI Q <GEL>
(Medicine for toothache)



Medicine for toothache

Class 2 drug

Konjisui Q is an EPC-formulated gel-type medication for toothache. Fast-acting for unbearable toothache, with long-lasting pain relief.

MADE IN JAPAN

Indications

toothache

Dosage and Administration

Apply an appropriate amount to a cotton bud or swab, and insert into the hole of the decayed tooth, or apply an appropriate amount directly over the hole of the decayed tooth.

(Precautions related to Dosage and Administration)

- (1) Please respect the stipulated application method.
- (2) Do not apply to areas other than the painful tooth cavity.
- (3) When used for children, please use under the guidance and supervision of the guardian.
- (4) This product is for dental use only. Do not use for ophthalmological or other uses.
- (5) This product will not harm the hard quality of the tooth (enamel and dentine quality); however, if excessive product spreads to areas other than the tooth, taste will temporarily change. Hence, please wipe off such excessive portion with a gauze.
- (6) If the product gets on the face or around the mouth by mistake, rinse immediately with water and wipe the product off.

※This product cannot cure the cause of toothache such as tooth decay, and so it is recommended to receive fundamental treatment from a dentist without delay.

Ingredient, content (per 100g), and action

Ingredient	Content	Action
eugenol	5.0g	Pain relief
dl-camphor	4.0g	
phenol	2.0g	Pain relief and bactericidal action.

Contains carboxyvinyl polymer, sodium hydroxide, fennel oil, and propylene glycol as additives.

Precautions

Things to avoid

(if not respected, the present symptoms will exacerbate, and side effects will readily develop)

Do not use the product in the following areas:

Gums, and lips

Things to consult

1. Before using the product, the following individuals should consult their doctor, dentist, pharmacist, or registered sales clerk:
 - (1) Individuals receiving medical treatment from a doctor or dentist.
 - (2) Individuals who have a history of drug-induced allergic symptoms.
2. After usage, if any of the following symptoms occur, they could be side effects; thus, discontinue usage of the product immediately, and please consult your doctor, pharmacist, or registered sales clerk with this document.

Site concerned	Symptoms
Skin	Eruption, redness, and itchiness

3. If symptoms do not improve after usage for 5 - 6 times, discontinue usage, and please consult your doctor, dentist, pharmacist, or registered sales clerk with this document.