



コウナキッズ

KOUNAKIZZU

Medicine for mouth ulcer and sore throat.

Class 3 drug

Spray to alleviate inflammation and pain of mouse ulcer.

Child-friendly grape taste

MADE IN JAPAN

Indications

Stomatitis, throat pain caused by throat inflammation, roughness of the throat, swelling of the throat, throat discomfort, and voice hoarseness

Dosage and Administration

Spray the affected area with an adequate amount several times a day. Furthermore, when using for the first time, press several times until the liquid medicine comes out.

(Precautions related to Dosage and Administration)

- (1) the liquid medicine will enter the bronchi and lungs if used while inhaling, and therefore, when using for the back of the throat, spray while making a vocalization.
 - (2) While using, make sure that the spray nozzle does not touch the inside of the mouth.
 - (3) Please strictly respect the application method and dosage.
 - (4) When used for children, please use under the guidance and supervision of the guardian.
 - (5) Keep out of eyes. If by chance the product gets in the eyes, rinse immediately with water or tepid water.
- Furthermore, when symptoms are severe, please receive medical attention from an ophthalmologist.

Ingredient, content (per 100g), and action

Ingredients	Content	Action
sodium gualenate hydrate	0.02g	Acts directly on the mucous membrane to alleviate inflammation, and improve symptoms such as pain and swelling.

Contains glycerin, sodium edetate, D-sorbitol, sodium saccharin, sodium hydrogen phosphate, sodium hydrogen carbonate, cetylpyridinium chloride hydrate, polyoxyethylene hardened castor oil, ethanol, and fragrance as additives.

Precautions

Things to consult

1. Before using the product, the following individuals should consult their doctor, dentist, pharmacist, or registered sales clerk:
 - (1) Individuals receiving medical treatment from a doctor or dentist.
 - (2) Individuals who have a history of drug-induced allergic symptoms.
 - (3) Individuals with severe inflammation in the mouth.
2. In the following instances, discontinue usage of the spray immediately, and please consult your doctor, pharmacist, or registered sales clerk with this document.

Site concerned	Symptoms
Mouth	Irritation
Skin	eruption, redness, and itchiness

3. When symptoms do not improve despite usage for 5 - 6 days.